Secondhand Marijuana Smoke

"Smoke is smoke. Both tobacco and marijuana smoke impair blood vessel function similarly. People should avoid both, and governments who are protecting people against secondhand smoke exposure should include marijuana in those rules."

-Matthew Springer, cardiovascular researcher and Associate Professor of Medicine, University of California, San Francisco

Facts about secondhand marijuana smoke:

- Marijuana smoke is created by burning components of plants in the genus Cannabis.
- Secondhand marijuana smoke is a complex chemical mixture of smoke emitted from combusted marijuana and the smoke that is exhaled by the user.
- Secondhand marijuana smoke contains fine particulate matter that can be breathed deeply into the lungs.
- Secondhand marijuana smoke contains many of the same cancer-causing substances and toxic chemicals as secondhand tobacco smoke. Some of the known carcinogens or toxins present in marijuana smoke include: acetaldehyde, ammonia arsenic, benzene, cadmium, chromium, formaldehyde, hydrogen cyanide, isoprene, lead, mercury, nickel, and quinoline.
- Marijuana smoke contains tetrahydrocannabinol (THC), the active chemical in cannabis.

Health risks of exposure to secondhand marijuana smoke:

Since marijuana is illegal under federal law, there have been a limited number of studies examining health risks associated with marijuana use and exposure in the United States. Health risks from primary and secondhand smoke exposure may also be difficult to determine as marijuana is often used in combination with tobacco.

However, peer-reviewed and published studies do indicate that exposure to secondhand marijuana smoke may have health and safety risks for the general public, especially due to its similar composition to secondhand tobacco smoke.

- Secondhand smoke from combusted marijuana contains fine particulate matter that can be breathed deeply into the lungs,ⁱⁱ which can cause lung irritation, asthma attacks, and makes respiratory infections more likely. Exposure to fine particulate matter can exacerbate health problems especially for people with respiratory conditions like asthma, bronchitis, or COPD.ⁱⁱⁱ
- Significant amounts of mercury, cadmium, nickel, lead, hydrogen cyanide, and chromium, as well as 3 times the amount of ammonia, are found in mainstream marijuana smoke than is in tobacco smoke.
- In 2009, the California Office of Environmental Health Hazard Assessment added marijuana smoke to its Proposition 65 list of carcinogens and reproductive toxins, also known as the Safe Drinking Water and Toxic Enforcement Act of 1986. It reported that at least 33 individual constituents present in both marijuana smoke and tobacco smoke are Proposition 65 carcinogens.^{v, vi}

- Secondhand smoke from marijuana has many of the same chemicals as smoke from tobacco, including those linked to lung cancer.^{vii}
- Secondhand marijuana exposure impairs blood vessel function. Published studies on rats show
 that thirty minutes of exposure to secondhand marijuana smoke at levels comparable to those
 found in restaurants that allow cigarette smoking led to substantial impairment of blood vessel
 function. Marijuana smoke exposure had a greater and longer-lasting effect on blood vessel
 function than exposure to secondhand tobacco smoke.
- One minute of exposure to marijuana SHS substantially impairs endothelial function in rats for at least 90 minutes, considerably longer than comparable impairment by tobacco SHS. The findings in rats suggest that SHS can exert similar adverse cardiovascular effects regardless of whether it is from tobacco or marijuana.
- Secondhand marijuana smoke and secondhand tobacco smoke is similar in many ways. More
 research is needed, but the current body of science shows that both tobacco and marijuana
 smoke have similar chemical composition and suggests that they may have harmful
 cardiovascular health effects, such as atherosclerosis (partially blocked arteries), heart attack,
 and stroke.^x
- Particle concentrations from dabbing and vaporizing cannabis can create levels of indoor air
 pollution similar as those seen in extreme air pollution events like wildfires and severe industrial
 pollution. Exposure at these concentrations can cause cardiovascular and respiratory disease.xi
- People who are exposed to secondhand marijuana smoke can have detectable levels of THC (tetrahydrocannabinol) in their blood and urine. xii
- Marijuana also can be contaminated with mold, insecticides or other chemicals that may be released in secondhand smoke.

Including Marijuana Smoking in Smokefree Public Place and Workplace Laws:

- Everyone has the right to breathe smokefree air. Smokefree policies are designed to protect the public and all workers from exposure to the health hazards caused by exposure to secondhand tobacco smoke. The same should be true for secondhand marijuana smoke.
- The percent of U.S. adults who use marijuana more than doubled from 4.1% to 9.5% between 2001-2002 and 2012-2013, xiv which may also indicate an increase in exposure to secondhand marijuana smoke.
- The American Society for Heating, Refrigeration, and Air Conditioning Engineering (ASHRAE) is the organization that develops engineering standards for building ventilation systems. ASHRAE now bases its ventilation standard for acceptable indoor air quality on an environment that is completely free from secondhand tobacco smoke, secondhand marijuana smoke, and emissions from electronic smoking devices.xv
- In order to protect public health, improve consistency, and aid enforcement, smokefree laws for
 public places and workplaces should include tobacco as well as marijuana, whether it is smoked
 or aerosolized. Allowing marijuana smoking in places where smoking is now prohibited could
 undermine laws that protect the public from exposure to secondhand smoke. The Tobacco
 Control Legal Consortium issued an informative brief on Lessons from Tobacco Control for Marijuana Regulation.

- Smokefree policies provide incentives to quit smoking, help denormalize smoking behavior, and
 are particularly effective among youth and young adults who are vulnerable to visual cues and
 social norms of smoking. It is likely that smokefree policies for marijuana will have a similar effect.
- As of January, 2021, 774 localities and 29 states/territories/commonwealths restrict marijuana
 use in some or all smokefree spaces. Of these, 417 localities and 17
 states/territories/commonwealths prohibit smoking and vaping of recreational and medical
 marijuana in one or more of the following venues: non-hospitality workplaces, restaurants, bars,
 and/or gambling facilities.

In the interest of public health, the use of combustible or aerosolized marijuana should be prohibited wherever tobacco smoking is prohibited.

ANR Foundation's Position on Exposure to Secondhand Marijuana Smoke:

Marijuana smoke is a form of indoor air pollution. Therefore, ANR, our lobbying organization, includes marijuana within the definition of smoking, and all of our model laws and policies include a prohibition on smoking marijuana wherever smoking of tobacco products is not allowed. Our organization does not have a position on whether marijuana should be legalized; we are committed to smokefree protections from secondhand smoke from tobacco products, marijuana and aerosol from electronic smoking devices.

Nobody should have to breathe secondhand marijuana smoke at work, in public, or where they live. If we want healthy, smokefree air for workers and the public, then products like marijuana and electronic smoking devices (which can be used to "vape" a wide range of substances, including marijuana and hash oil) must not be used in smokefree environments where others are forced to breathe the secondhand emissions.

References

¹ Moir, D., et al., A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. Chem Res Toxicol 21: 494-502. (2008). http://www.ncbi.nlm.nih.gov/pubmed/18062674

¹ Hillier, FC.; et al. "Concentration and particle size distribution in smoke from marijuana cigarettes with different Δ9-tetrahydrocannabinol content." Fundamental and Applied Toxicology. Volume 4, Issue 3, Part 1, June 1984, Pages 451-454. http://www.sciencedirect.com/science/article/pii/0272059084902021

[&]quot;Air and Health: Particulate Matter." National Environmental Public Health Tracking Network, U. S. Environmental Protection Agency. http://ephtracking.cdc.gov/showAirHealth.action#ParticulateMatter

iv Moir, D., et al., A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. Chem Res Toxicol 21: 494-502. (2008). http://www.ncbi.nlm.nih.gov/pubmed/18062674
v "Evidence on the Carcinogenicity of Marijuana Smoke." Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. August 2009. http://oehha.ca.gov/prop65/hazard ident/pdf zip/FinalMJsmokeHID.pdf

Wang, X., et al., "Brief exposure to marijuana secondhand smoke impairs vascular endothelial function" (conference abstract). *Circulation* 2014; 130: A19538. http://circ.ahajournals.org/content/130/Suppl_2/A19538.abstract

vii "Evidence on the Carcinogenicity of Marijuana Smoke." Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. August 2009.

http://oehha.ca.gov/prop65/hazard_ident/pdf_zip/FinalMJsmokeHID.pdf viii Wang, X., et al., "Brief exposure to marijuana secondhand smoke impairs vascular endothelial function" (conference abstract). Circulation 2014; 130: A19538. http://circ.ahajournals.org/content/130/Suppl_2/A19538.abstract

Wang, X.; Derakhshandeh, R.; Liu, J.; Narayan, S.; Nabavizadeh, P.; Le, S.; Danforth, O.M.; Pinnamaneni, K.; Rodriguez, H.J.; Luu, E.; Sievers, R.E.; Schick, S.F.; Glantz, S.A.; Springer, M.L., "One minute of marijuana secondhand smoke exposure

substantially impairs vascular endothelial function," *Journal of the American Heart Association* 5(8): e003858, August 2016. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5015303/

https://tobacco.ucsf.edu/sites/tobacco.ucsf.edu/files/u9/MSHS%20fact%20sheet%20for%20CA%204-13-15.pdf

- ^{xii} Herrmann ES, et al., "Non-smoker exposure to secondhand cannabis smoke II: Effect of room ventilation on the physiological, subjective, and behavioral/cognitive effects." *Drug and Alcohol Dependence*. 2015 Jun 1;151:194-202. http://www.ncbi.nlm.nih.gov/pubmed/25957157
- <u>UCDavis Researchers, research letter published online in the journal of Clinical Microbiology and Infection,</u> "A microbiome assessment of medical marijuana" April 2017Volume 23, Issue 4, Pages 269–270 https://www.clinicalmicrobiologyandinfection.com/article/S1198-743X(16)30605-X/pdf
- ^{xiv} Hasin DS, et al. "Prevalence of Marijuana Use Disorders in the United States Between 2001-2002 and 2012-2013." JAMA Psychiatry. Published online October 21, 2015. http://archpsyc.jamanetwork.com/article.aspx?articleid=2464591
- ^{xv} ANSI/ASHRAE Standard 62.1-2016 Ventilation for Acceptable Indoor Air Quality. Atlanta, GA: American Society of Heating, Refrigerating, and Air-Conditioning Engineers, Inc.

https://www.ashrae.org/technical-resources/standards-and-guidelines/read-only-versions-of-ashrae-standards

xvi Cork, Kerry. "Toking, Smoking & Public Health: Lessons from Tobacco Control for Marijuana Regulation." Tobacco Control Legal Consortium. June 2015. http://www.publichealthlawcenter.org/sites/default/files/resources/tclc-synopsis-marijuana-tobacco-2015 0.pdf

For more information, visit https://nonsmokersrights.org/marijuana-smoke or call us at 510-841-3032.

May be reprinted with appropriate attribution to the American Nonsmokers' Rights Foundation, © 2021

2021-02 [FS-45]

^x Springer, M.L.; Glantz, S.A." Marijuana Use and Heart Disease: Potential Effects of Public Exposure to Smoke," University of California at San Francisco. April 13, 2015.

^{xi} Jaques, P, Zalay, M, Huang, A, Jee, K, Schick, SF "Measuring Aerosol Particle Emissions from Cannabis Vaporization and Dabbing", Proceedings of the 15th Meeting of the International Society for Indoor Air Quality and Climate. July 22-27, 2018. Philadelphia, PA.